

DAFTAR PUSTAKA

- [1] M. M. Addi, N. A. Z. Abidin, and S. A. Daud, “Development of a portable phototherapy garment (PPG) for jaundice treatment,” *IECBES 2016 - IEEE-EMBS Conf. Biomed. Eng. Sci.*, pp. 405–410, 2016, doi: 10.1109/IECBES.2016.7843482.
- [2] V. K. Bhutani *et al.*, “Phototherapy to prevent severe neonatal hyperbilirubinemia in the newborn infant 35 or more weeks of gestation,” *Pediatrics*, vol. 128, no. 4, 2011, doi: 10.1542/peds.2011-1494.
- [3] A. K. Surya Dewi, I. M. Kardana, and K. Suarta, “Efektivitas Fototerapi Terhadap Penurunan Kadar Bilirubin Total pada Hiperbilirubinemia Neonatal di RSUP Sanglah,” *Sari Pediatr.*, vol. 18, no. 2, p. 81, 2016, doi: 10.14238/sp18.2.2016.81-6.
- [4] R. Pepenene, R. Sieberhagen, and N. Nel-Sakharova, “Radiometry of Phototherapy (Blue Light Therapy) in South Africa,” no. September 2014, 2011, doi: 10.13140/2.1.3972.7360.

- [5] F. P. Facchini, “Standardizing the calibration of phototherapy devices,” *J. Pediatr. (Rio. J.)*, vol. 77, no. 2, pp. 67–74, 2001, doi: 10.2223/jped.179.
- [6] M. T. A. Sampurna *et al.*, “Current phototherapy practice on Java, Indonesia,” *BMC Pediatr.*, vol. 19, no. 1, pp. 1–9, 2019, doi: 10.1186/s12887-019-1552-1.
- [7] D. Bunyaniah, “Pengaruh Fototerapi Terhadap Derajat Ikterik Pada Bayi Baru Lahir Di Rsud Dr. Moewardi Surakarta,” pp. 1–21, 2013.
- [8] P. D. M. Jeffrey Maisels, M.B., B.Ch., and Antony F. McDonagh and This, “Phototherapy for neonatal jaundice,” *N. Engl. J. Med.*, vol. 358, no. 23, pp. 2523–2524, 2008.
- [9] Permenkes RI, “Permenkes R1 No 54 Tahun 2015,” 2015.
- [10] H. Moseley *et al.*, “Guidelines on the measurement of ultraviolet radiation levels in ultraviolet phototherapy: Report issued by the British Association of Dermatologists and British

Photodermatology Group 2015,” *Br. J. Dermatol.*, vol. 173, no. 2, pp. 333–350, 2015, doi: 10.1111/bjd.13937.

- [11] H. J. Vreman, R. J. Wong, and D. K. Stevenson, “Phototherapy: Current methods and future directions,” *Semin. Perinatol.*, vol. 28, no. 5, pp. 326–333, 2004, doi: 10.1053/j.semperi.2004.09.003.
- [12] J. Webster, R. Blythe, and F. Nugent, “An appraisal of the use of the Kramer’s scale in predicting hyperbilirubinaemia in healthy full term infants.,” *Birth Issues*, vol. 14, no. 3, pp. 83–89, 2006.
- [13] D. Kr and V. Artemov, “(12) Patent Application Publication (10) Pub. No.: US 2012/0184582 A1,” US 2012/0280114A1, 2012.
- [14] S. M. Reda, K. A. Mohammad, and S. M. El-faramawy, “Construction and characterization of a phototherapy radiometer for optical radiation measurements,” *Indian J. Pure Appl. Phys.*, vol.

56, no. 5, pp. 379–382, 2018.

- [15] T. R. Ganjar Heru Purnomo, Bedjo Utomo, “Phototeraphy Radiometer Dengan Penyimpanan Data Pengukuran Pada SDCard,” p. 7, 2018.
- [16] D. M. Clarkson and P. Satodia, “Use of a hand-held spectroradiometer for the measurement of neonatal phototherapy lamp outputs,” *Med. Eng. Phys.*, vol. 73, pp. 107–111, 2019, doi: 10.1016/j.medengphy.2019.07.001.
- [17] Ichwan Syahrul Bahtiar, Andjar Pudji, and I Dewa Gede Hari Wisana, “Phototherapy Radiometer with AS7262 Sensor,” *J. Electron. Electromed. Eng. Med. Informatics*, vol. 1, no. 1, pp. 39–45, 2019, doi: 10.35882/jeeemi.v1i1.8.
- [18] S. Mathindas, R. Wilar, and A. Wahani, “Hiperbilirubinemia Pada Neonatus,” *J. Biomedik*, vol. 5, no. 1, 2013, doi: 10.35790/jbm.5.1.2013.2599.
- [19] S. Pratesi, S. Di Fabio, C. Bresci, C. Di Natale, S. Bar, and C. Dani, “Broad-Spectrum Light versus

Blue Light for Phototherapy in Neonatal Hyperbilirubinemia: A Randomized Controlled Trial,” *Am. J. Perinatol.*, vol. 32, no. 8, pp. 779–784, 2014, doi: 10.1055/s-0034-1396685.

- [20] O. F. Pediatrics, N. Infant, and M. Weeks, “Management of Hyperbilirubinemia in the Newborn Infant 35 or More Weeks of Gestation,” *Pediatrics*, vol. 114, no. 1, pp. 297–316, 2004, doi: 10.1542/peds.114.1.297.
- [21] F. Pineda-Lopez *et al.*, “Light blue led for bilirubin treatment in newborns: Automatic phototherapy prototype,” *Proc. 2017 IEEE 24th Int. Congr. Electron. Electr. Eng. Comput. INTERCON 2017*, pp. 0–3, 2017, doi: 10.1109/INTERCON.2017.8079715.
- [22] M. S. Kosim, R. Soetandio, and M. Sakundarno, “Dampak Lama Fototerapi Terhadap Penurunan Kadar Bilirubin Total pada Hiperbilirubinemia Neonatal,” *Sari Pediatr.*, vol. 10, no. 3, p. 201, 2016, doi: 10.14238/sp10.3.2008.201-6.

- [23] G. M. PRAVEEN KUMAR, SRINIVAS MURKI*, “Light-Emitting Diodes versus Compact Fluorescent Tubes for Phototherapy.,” *Indian pediatrics*, vol. 47, no. 11. p. 979, 2010.