

DAFTAR PUSTAKA

- Anxiety, D. *et al.* (2016) *Nursing Diagnosis Handbook*.
- Azizah, N. U. R. (2020) 'SCOPING REVIEW : PENGARUH POSISI PRONE TERHADAP SATURASI OKSIGEN (SpO₂) PADA PASIEN COVID-19 FAKULTAS KEDOKTERAN SCOPING REVIEW: PENGARUH POSISI PRONE TERHADAP SATURASI OKSIGEN (SpO₂) PADA PASIEN COVID-19'.
- Budiono (2016) *Konsep Dasar Keperawatan*. edisi 1. Edited by A. B. Darmanto and A. Sutisna. Jakarta Selatan: Pusdik SDM Kesehatan.
- Cohen, D. *et al.* (2020) 'Beneficial effect of awake prone position in hypoxaemic patients with COVID-19: case reports and literature review', *Internal Medicine Journal*, 50, pp. 997–1000. doi: 10.1111/imj.14926.
- Coronavirus disease (COVID-19)* (no date). Available at: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/question-and-answers-hub/q-a-detail/coronavirus-disease-covid-19#:~:text=symptoms> (Accessed: 7 August 2021).
- Diah, P. (2021) 'View of EFEKTIFITAS POSISI PRONASI PADA PASIEN COVID-19 DENGAN GANGGUAN PERNAPASAN TANPA INTUBASI DI RUANG ICU : LITERATUR REVIEW', *Jurnal Ilmiah Keperawatan (Scientific Journal of Nursing)*, 7(7), p. 2. Available at: <http://www.journal.stikespemkabjombang.ac.id/index.php/jikep/article/view/845/598> (Accessed: 18 October 2021).
- Doussot, A. *et al.* (2020) 'Prone Positioning for Severe Acute Respiratory Distress Syndrome in COVID-19 Patients by a Dedicated Team A Safe and Pragmatic Reallocation of Medical and Surgical Work Force in Response to the Outbreak'. doi: 10.1097/SLA.0000000000004265.
- Dubosh, N. M. *et al.* (2021) 'Early, awake proning in emergency department patients with COVID-19', *The American Journal of Emergency Medicine*. W.B. Saunders, 46, pp. 640–645. doi: 10.1016/J.AJEM.2020.11.074.
- Elharrar, X. *et al.* (2020) 'Use of Prone Positioning in Nonintubated Patients With COVID-19 and Hypoxemic Acute Respiratory Failure', *JAMA*. American Medical Association, 323(22), p. 2336. doi: 10.1001/JAMA.2020.8255.
- Fatima, A. *et al.* (2020) 'Literature Review of Covid-19: Virology, Pathogenesis, and Clinical Manifestations.', *Human Relations*, 3(1), pp. 1–8. Available at: <http://search.ebscohost.com/login.aspx?direct=true&AuthType=ip,shib&db=bth&AN=92948285&site=eds->

live&scope=site%0Ahttp://bimpactassessment.net/sites/all/themes/bcorp_impact/pdfs/em_stakeholder_engagement.pdf%0Ahttps://www.globebus.com/help/helpFiles/CDJ-Pa.

- Huang, C. *et al.* (2020) ‘Clinical features of patients infected with 1. C H, Y W, X L, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* (London, England). 2020;395(10223):497-506. doi:10.1016/S0140-6736(20)30183-5 2019 novel coronav’, *Lancet* (London, England). *Lancet*, 395(10223), pp. 497–506. doi: 10.1016/S0140-6736(20)30183-5.
- Isbaniah, F. (2020) ‘Pedoman Pencegahan dan Pengendalian Coronavirus Disease (COVID-19)’, *Germas*, pp. 0–115. Available at: https://infeksiemerging.kemkes.go.id/download/REV-04_Pedoman_P2_COVID-19__27_Maret2020_TTD1.pdf [Diakses 11 Juni 2021].
- Jochmans, S. *et al.* (2020) ‘Duration of prone position sessions: a prospective cohort study’, *Ann. Intensive Care*, 10, p. 66. doi: 10.1186/s13613-020-00683-7.
- Kementerian Kesehatan Republik Indonesia (2020) ‘Situasi Terkini Perkembangan (COVID-19)’, *Kemendes*, (September), pp. 17–19. Available at: https://covid19.kemkes.go.id/download/Situasi_Terkini_050520.pdf.
- Kuriasih, D. N. *et al.* (2021) ‘Pengaruh prone position terhadap peningkatan oksigenasi pada pasien covid-19: Systematic review’, *Holistik Jurnal Kesehatan*, 15(2), pp. 274–286. Available at: <http://ejournalmalahayati.ac.id/index.php/holistik/article/view/4894> (Accessed: 25 October 2021).
- Levani, Prastya and Mawaddatunnadila (2021) ‘Coronavirus Disease 2019 (COVID-19): Patogenesis, Manifestasi Klinis dan Pilihan Terapi’, *Jurnal Kedokteran dan Kesehatan*, 17(1), pp. 44–57. Available at: <https://jurnal.umj.ac.id/index.php/JKK/article/view/6340>.
- Li, Q. *et al.* (2020) ‘Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia’, *New England Journal of Medicine*, 382(13), pp. 1199–1207. doi: 10.1056/nejmoa2001316.
- Nyoman, N., Darmayanti, T. and Oktamianti, P. (2019) ‘Analisis Kompetensi Perawat Ruang Intensif (Intensive Care Unit)’, *Jurnal Sumber Daya Manusia Kesehatan*, 1, p. 1.
- PDPI (2020) ‘RESPIROLOGI INDONESIA’, *CPD Infection*, 3(1), pp. 9–12.
- PPNI (2016) *SDKI – Standart Diagnosis Keperawatan Indonesia*. Jakarta. Available at: <http://snars.web.id/sdki/d0005-pola-napas-tidak-efektif/>

(Accessed: 22 August 2021).

- Sukmana, M. and Yuniarti, F. A. (2020) 'The Pathogenesis Characteristics and Symptom of Covid-19 in the Context of Establishing a Nursing Diagnosis', *Jurnal Kesehatan Pasak Bumi Kalimantan*, 3(1), p. 21. doi: 10.30872/j.kes.pasmi.kal.v3i1.3748.
- Susilo, A. *et al.* (2020) 'Coronavirus Disease 2019: Tinjauan Literatur Terkini', *Jurnal Penyakit Dalam Indonesia*, 7(1), p. 45. doi: 10.7454/jpdi.v7i1.415.
- Vollenberg, R. *et al.* (2021) 'Clinical Medicine Prone Position in Mechanically Ventilated COVID-19 Patients: A Multicenter Study Prone Position in Mechanically'. doi: 10.3390/jcm10051046.
- Weiss, T. T. *et al.* (2021) 'Prone positioning for patients intubated for severe acute respiratory distress syndrome (ARDS) secondary to COVID-19: a retrospective observational cohort study', *British Journal of Anaesthesia*. Elsevier, 126(1), pp. 48–55. doi: 10.1016/J.BJA.2020.09.042.
- WHO (2020) 'Transmisi SARS-CoV-2: implikasi terhadap kewaspadaan pencegahan infeksi', pp. 1–10.
- Widysanto, A. *et al.* (2020) 'Happy hypoxia in critical COVID-19 patient: A case report in Tangerang, Indonesia', *Physiological Reports*. Wiley-Blackwell, 8(20). doi: 10.14814/PHY2.14619.