

DAFTAR PUSTAKA

- Adcock Funk, D. M., Lippi, G., & Favaloro, E. J. (2012). Quality standards for sample processing, transportation, and storage in hemostasis testing. *Seminars in Thrombosis and Hemostasis*, 38(6), 576–585. <https://doi.org/10.1055/s-0032-1319768>
- Alesci, S., Borggrefe, M., & Dempfle, C. E. (2009). Effect of freezing method and storage at - 20 °C and - 70 °C on prothrombin time, aPTT and plasma fibrinogen levels. *Thrombosis Research*, 124(1), 121–126. <https://doi.org/10.1016/j.thromres.2008.11.010>
- Anandani, G. M., & Parikh, S. B. (2018). Effect of pre-analytic variables on prothrombin time and activated partial thromboplastin time. *Journal of Clinical and Diagnostic Research*, 12(7), EC01–EC05. <https://doi.org/10.7860/JCDR/2018/32666.11719>
- Antovic, J. P., Blombäck, M., & Wiley, J. (2013). *Essential Guide to Blood Coagulation 2 nd Edition Edited by. www.wiley.com/wiley-blackwell.*
- Bennett, S. T., Rodgers, G. M., & Lehman, C. M. (2015). Hemostasis Screening Assays. In *Laboratory Hemostasis*. https://doi.org/10.1007/978-3-319-08924-9_5
- CLSI. (2008). Collection , Transport, and Processing of Blood Specimens for Testing Plasma-Based Coagulation Assays and Molecular Hemostasis Assays; CLSI Approved Guideline-Fifth Edition. H21-A5. *Nccls*, 28(5), H21-A5.

- Denessen, E. J. S., Jeurissen, M. L. J., Pereboom, R. M. T. A., Verhezen, P. W. M., & Henskens, Y. M. C. (2020). Determining the maximal storage time of centrifuged citrated samples for performing add-on routine coagulation tests. *Thrombosis Research*, *196*(July), 54–62. <https://doi.org/10.1016/j.thromres.2020.08.003>
- Durachim, A., & Astuti, D. (2018). *Bahan Ajar Teknologi Laboratorium Medik - Hemostasis* (Edisi Tahu). Badan Pengembangan dan Pembedayaan Sumber Daya Manusia Kesehatan Edisi Tahun 2018. http://bppsdmk.kemkes.go.id/pusdiksdmk/wp-content/uploads/2018/09/Hemostasis_SC.pdf
- Favaloro, E. J., Rcpa, F., Funk, D. M. A., & Lippi, G. (2018). *Pre-analytical Variables in Coagulation Testing*. *43*(2), 1–10. <https://doi.org/10.1309/LM749BQETKYPYPVM>
- Feng, L., Zhao, Y., Zhao, H., & Shao, Z. (2014). Effects of storage time and temperature on coagulation tests and factors in fresh plasma. *Scientific Reports*, *4*, 6–10. <https://doi.org/10.1038/srep03868>
- Foshat, M., Bates, S., Russo, W., Huerta, A., Albright, K., Giddings, K., Indrikovs, A., & Qian, Y. W. (2015). Effect of freezing plasma at -20°C for 2 weeks on prothrombin time, activated partial thromboplastin time, dilute Russell viper venom time, activated protein c resistance, and d-dimer levels. *Clinical and Applied Thrombosis/Hemostasis*, *21*(1), 41–47. <https://doi.org/10.1177/1076029614535973>

- Geelani, S., Wani, G. S., Khan, S. P., Qadri, S. M., Rasool, J., Quadri, S. S., & Khan, F. P. (2018). Effect of storage time on prothrombin time and activated partial thromboplastin time: study at a tertiary care center in Kashmir valley. *International Journal of Scientific Reports*, 4(7), 182. <https://doi.org/10.18203/issn.2454-2156.intjsciirep20182729>
- Hassan, N. A., Ali, E. W., Mohammed, N. Al., Ibrahim, I. K., Khartoum, & Sudan. (2013). The Effect of Freezing Storage of Citrated Plasma on Prothrombin Time and Activated Partial Thromboplastin Time. *Asin Journal of Biomedical and Pharmaceutical Sciences*, 2–3.
- Kemkes-Matthes, B., Fischer, R., & Peetz, D. (2011). Influence of 8 and 24-h storage of whole blood at ambient temperature on prothrombin time, activated partial thromboplastin time, fibrinogen, thrombin time, antithrombin and D-dimer. *Blood Coagulation and Fibrinolysis*, 22(3), 215–220. <https://doi.org/10.1097/MBC.0b013e328343f8bf>
- Keohane, E. M., Otto, C. N., & Walenga, J. M. (2019). *Rodak's Hematology: Clinical Principles and Applications* (Sixth Edit). Elsevier Health Sciences.
- Kitchen, S., Adcock, D. M., Dauer, R., Kristoffersen, A. H., Lippi, G., Mackie, I., Marlar, R. A., & Nair, S. (2021a). International Council for Standardisation in Haematology (ICSH) recommendations for collection of blood samples for coagulation testing. *International Journal of Laboratory Hematology*, 43(4), 571–580. <https://doi.org/10.1111/ijlh.13584>
- Kitchen, S., Adcock, D. M., Dauer, R., Kristoffersen, A. H., Lippi, G., Mackie, I.,

- Marlar, R. A., & Nair, S. (2021b). International Council for Standardization in Haematology (ICSH) recommendations for processing of blood samples for coagulation testing. *International Journal of Laboratory Hematology*, 43(6), 1272–1283. <https://doi.org/10.1111/ijlh.13702>
- Lippi, G., Salvagno, G., Montagnana, M., Lima-Oliveira, G., Guidi, G., & Favaloro, E. J. (2012). Quality standards for sample collection in coagulation testing. *Seminars in Thrombosis and Hemostasis*, 38(6), 565–575. <https://doi.org/10.1055/s-0032-1315961>
- Magnette, A., Chatelain, M., Chatelain, B., Ten Cate, H., & Mullier, F. (2016). Pre-analytical issues in the haemostasis laboratory: Guidance for the clinical laboratories. *Thrombosis Journal*, 14(1), 1–14. <https://doi.org/10.1186/s12959-016-0123-z>
- Marder, V. J., Aird, W. C., Bennett, J. S., Schulman, S., & White, G. C. (2012). *Hemostasis and Thrombosis: Basic Principles and Clinical Practice* (V. J. Marder, W. C. Aird, J. S. Bennett, S. Schulman, & G. C. White (eds.); Sixth). Wolters Kluwer Health.
- Musa, A., Kwaifa, I., Gusau, Y., Hassan, A.-A., & Ndakotsu, M. (2013). Effect of plasma storage on prothrombin time and activated partial thromboplastin time at a Nigerian public laboratory. *Sahel Medical Journal*, 16(1), 1. <https://doi.org/10.4103/1118-8561.112038>
- Rimac, V., & Coen Herak, D. (2017). Is it acceptable to use coagulation plasma samples stored at room temperature and 4°C for 24 hours for additional

prothrombin time, activated partial thromboplastin time, fibrinogen, antithrombin, and D-dimer testing? *International Journal of Laboratory Hematology*, 39(5), 475–481. <https://doi.org/10.1111/ijlh.12664>

Siregar, M. T., Wulan, W. S., Setiawan, D., & Nuryati, A. (2018). *Bahan Ajar Teknologi Laboratorium Medik - Kendali Mutu* (Edisi 2018). Badan Pengembangan dan Pemberdayaan Sumber Daya Manusia Kesehatan, Kementerian Kesehatan Republik Indonesia. http://bppsdmk.kemkes.go.id/pusdiksdmk/wp-content/uploads/2018/09/Kendali-Mutu_SC.pdf

Van Den Besselaar, A. M. H. P., Witteveen, E., & Van Der Meer, F. J. M. (2013). Long-term stability of frozen pooled plasmas stored at - 70 C, - 40 C, and - 20 C for prothrombin time and International Normalized Ratio (INR) assessment. *Thrombosis Research*, 131(4), 349–351. <https://doi.org/10.1016/j.thromres.2013.01.020>

Zhao, Y., & Lv, G. (2013). Influence of temperature and storage duration on measurement of activated partial thromboplastin time, D-dimers, fibrinogen, prothrombin time and thrombin time, in citrate-anticoagulated whole blood specimens. *International Journal of Laboratory Hematology*, 35(5), 566–570. <https://doi.org/10.1111/ijlh.12113>