

## DAFTAR PUSTAKA

- [1] S. R. Nadhiroh, “Faktor Yang Berhubungan Dengan Kejadian Stunting Pada Balita,” 2010.
- [2] N. F. Krebs *et al.*, “Meat consumption is associated with less stunting among toddlers in four diverse low-income settings,” *Food Nutr. Bull.*, vol. 32, no. 3, pp. 185–191, 2011.
- [3] UNCEF, “Key Facts and Figures on Nutrition,” vol. 140, no. April, pp. 3–6, 2013.
- [4] A. J. Prendergast and J. H. Humphrey, “The stunting syndrome in developing countries,” *Paediatr. Int. Child Health*, vol. 34, no. 4, pp. 250–265, 2014.
- [5] F. O. Aridiyah, N. Rohmawati, and M. Ririanty, “Faktor-faktor yang Mempengaruhi Kejadian Stunting pada Anak Balita di Wilayah Pedesaan dan Perkotaan ( The Factors Affecting Stunting on Toddlers in Rural and Urban Areas ),” vol. 3, no. 1, 2015.
- [6] Mega Orina Fitri, “Aplikasi Monitoring Perkembangan Status Gizi Anak Dan Balita Secara

Digital Dengan Metode Antropometri Berbasis Android,” vol. 2, no. April, pp. 140–149, 2017.

- [7] J. Sahar, R. Tutik, and S. Haryati, “Balita,” no. 1995, pp. 131–135, 2003.
- [8] T. Pustaka, “Hubungan Antara Keaktifan Ibu Dalam Kegiatan Posyandu dan Pola Makan Balita di Kelurahan Rajabasa Raya Kecamatan Rajabasa Kota Bandar Lampung,” pp. 7–43, 2010.
- [9] D. Teori, “Pengukuran Antropometri Dasar Teori,” no. 1966, 2001.
- [10] N. K. Sutiari, *Anthropometry dan Dietary Assessment*. 2017.
- [11] B. S. Zemel, E. M. Riley, and V. A. Stallings, “Evaluation Of Methodology For Nutritional Assessment In Children : Anthropometry, Body Composition, and Energy Expenditure,” *Annu. Rev. Nutr.*, vol. 17, no. 1, pp. 211–235, 1997.
- [12] Nathan A, Scobell A, *J. Chem. Inf. Model.*, vol. 53, no. 9, pp. 1689–1699, 2013.
- [13] A. Sen Kilic and Z. Öndogan, “Evaluating and defining body measurements according to anthropometric measurement technique for lower garment pattern preparation,” *Tekst. ve*

*Konfeksiyon*, vol. 26, no. 1, pp. 109–116, 2016.

- [14] M. A. McDowell, C. D. Fryar, C. L. Ogden, and K. M. Flegal, “Anthropometric reference data for children and adults: United States, 2003-2006,” *Natl. Health Stat. Report.*, no. 10, pp. 2003–2006, 2008.
- [15] C. V. Dezenberg, T. R. Nagy, B. A. Gower, R. Johnson, and M. I. Goran, “Predicting body composition from anthropometry in pre-adolescent children,” *Int. J. Obes.*, vol. 23, no. 3, pp. 253–259, 1999.
- [16] Kementerian Kesehatan RI, “Buku Saku Nasional Pemantauan Status Gizi 2017,” 2017.
- [17] A. Wulan, “Hubungan Pengetahuan Ibu tentang Gizi balita dengan Kejadian Stunting Anak,” *Univ. Muhammadiyah Surakarta*, no. 2017, pp. 5–18, 2010.
- [18] M. De Onis, M. Blössner, and E. Borghi, “Prevalence and trends of stunting among pre-school children, 1990-2020,” *Public Health Nutr.*, vol. 15, no. 1, pp. 142–148, 2012.
- [19] M. de Onis and F. Branca, “Childhood stunting: A global perspective,” *Matern. Child Nutr.*, vol. 12,

pp. 12–26, 2016.

- [20] I. Guide, “Interpretation Guide,” *Nutr. Landacape Inf. Syst.*, pp. 1–51, 2012.
- [21] N. K. Aryastami, A. Shankar, N. Kusumawardani, B. Besral, A. B. Jahari, and E. Achadi, “Low birth weight was the most dominant predictor associated with stunting among children aged 12–23 months in Indonesia,” *BMC Nutr.*, vol. 3, no. 1, pp. 1–6, 2017.
- [22] J. Hoddinott, H. Alderman, J. R. Behrman, L. Haddad, and S. Horton, “The economic rationale for investing in stunting reduction,” *Matern. Child Nutr.*, vol. 9, no. S2, pp. 69–82, 2013.
- [23] K. G. Dewey and K. Begum, “Long-term consequences of stunting in early life,” *Matern. Child Nutr.*, vol. 7, no. SUPPL. 3, pp. 5–18, 2011.
- [24] Fauziah L, Rahman N, Hermiyanti, “Faktor Risiko Kejadian Gizi Kurang pada Balita Usia 24-59 Bulan Di Kelurahan Taipa Kota Palu,” (2017) 4 (3) 29-59.
- [25] E. Waliyo, “Gizi Kurus (Wasting) Pada Balita,” pp. 1–7, 2013.
- [26] R. Afriyani and N. Malahayati, “Faktor-Faktor Yang Mempengaruhi Kejadian Wasting Pada Balita Usia 1-5 Tahun,” pp. 66–72, 2015.

