

## DAFTAR PUSTAKA

- Anush, M. M., Ashok, V. K., Sarma, R. I. N., & Pillai, S. K. (2019). Role of c-reactive protein as an indicator for determining the outcome of sepsis. *Indian Journal of Critical Care Medicine*, 23(1). DOI: 10.5005/jp-journals-10071-23105
- Bansal, C., Agrawal, R., Sukumaran, T. (2013). In *IAP Textbook of Pediatrics*. Retrieved from <https://doi.org/10.5005/jp/books/11894>
- Bapennas. (2018). *Tujuan SDGs Ke-3 Kehidupan Sehat dan Sejahtera*. Bapennas.
- Bayana, E., Endale, K., Olani, A., Biratu, Y., Tegenu, K. (2020). 'Neonatal Sepsis among Neonates at Public Hospitals in Jimma, Ethiopia', *International Journal of Pediatrics*, 8(9), pp. 12011-12021. doi: 10.22038/ijp.2020.44435.3681
- Borghesi, A., et al. (2014). IS-013 Neonatal Sepsis, New Preventive Strategies. *Archives of Disease in Childhood*, 99(Suppl 2). Retrieved from <http://dx.doi.org/10.1136/archdischild-2014-307384.13>
- Camacho-Gonzalez, A., Spearman, P. W., & Stoll, B. J. (2013). Neonatal Infectious Diseases. Evaluation of Neonatal Sepsis. In *Pediatric Clinics of North America (Vol. 60, Issue 2)*. doi: 10.1016/j.pcl.2012.12.003 .
- Celik, I. H., et al. (2022). Diagnosis of neonatal sepsis: the past, present and future. *Pediatric research*, 91(2), 337–350. DOI: 10.1038/s41390-021-01696-z
- Chandra, H. K., Fatoni, A. Z., (2021). Peranan C-Reactive Protein (CRP) pada Pasien Sepsis di Intensive Care Unit (ICU). *Journal of Anaesthesia and Pain*. DOI: <http://dx.doi.org/10.21776/ub.jap.2021.002.01.01>
- Chaurasiya OS, Ahmad T, Chhabra K, Nath D, Shobhne HJ. (2017). Procalcitonin Level in Neonatal Sepsis. *PJSR* ; 10(1):29-32.
- Dewi, J. (2018). "Peran Procalcitonin sebagai Marker Infeksi." *Cermin Dunia Kedokteran*, vol. 45, no. 7. doi:10.55175/cdk.v45i7.649.
- Emad A. Morad, Rehab A. Rabie, Mohamed A. Almalky, Manar G. Gebriel. (2020). "Evaluation of Procalcitonin, C-Reactive Protein, and Interleukin-6 as Early Markers for Diagnosis of Neonatal Sepsis", *International Journal of Microbiology*, vol. 2020, Article ID 8889086, 9 pages. doi: 10.1155/2020/8889086
- F. Hasan, S. A. Khan, M. K. Maharroof, and N. Muhammed. (2017). "Role of procalcitonin in early diagnosis of neonatal sepsis," *International Journal of Contemporary Pediatrics*, 4(2):7. doi: 10.18203/2349-3291.ijcp20170477

- Geni, L., & Panjaitan, L. M. R. (2019). Hubungan Kadar Procalcitonin (PCT) dengan C-Reactive Protein (Crp) Pada Pasien Infeksi Di Rumah Sakit Pluit. *Anakes : Jurnal Ilmiah Analis Kesehatan*, 5(1). Retrieved from <https://doi.org/10.37012/anakes.v5i1.333>
- Ikatan Dokter Anak Indonesia. (2016). Konsensus Diagnosis dan Tata Laksana Sepsis pada Anak. *Badan Penerbit Ikatan Dokter Anak Indonesia*.
- Iskandar, A. U. (2015) 'Pengambilan Sampel Darah Universitas Muhammadiyah Semarang'.
- Kemkes RI. (2021). KEPUTUSAN MENTERI KESEHATAN REPUBLIK INDONESIA NOMOR HK.01.07/MENKES/4722/2021 TENTANG PEDOMAN NASIONAL PELAYANAN KEDOKTERAN TATA LAKSANA SEPSIS PADA ANAK. *Jdih.Kemkes.Go.Id*.
- Kiriakopoulos, A., Giannakis, P., & Menenakos, E. (2022). Calcitonin: current concepts and differential diagnosis. *Therapeutic advances in endocrinology and metabolism*, 13. doi: 10.1177/20420188221099344
- Levinson T, Wasserman A. (2022). C-Reactive Protein Velocity (CRPv) as a New Biomarker for the Early Detection of Acute Infection/Inflammation. *Int J Mol Sci*. 23(15):8100. doi: 10.3390/ijms23158100.
- Markic, J., Saraga, M., & Dahlem, P. (2017). Sepsis Biomarkers in Neonates and Children: C-Reactive Protein and Procalcitonin. In *Journal of Child Science (Vol. 7, Issue 1)*. DOI:10.1055/s-0037-1603895.
- Menteri Kesehatan Republik Indonesia. (2017). Keputusan menteri kesehatan republik indonesia nomor HK.01.07/MENKES/342/2017 tentang pedoman nasional pelayanan kedokteran tata laksana sepsis. Jakarta: Menkes RI.
- Murthy, S., et.al. (2019). Risk factors of neonatal sepsis in India: A systematic review and meta-analysis. In *PLoS ONE* (Vol. 14, Issue 4). <https://doi.org/10.1371/journal.pone.0215683>
- Procianoy, R. S., & Silveira, R. C. (2020). The challenges of neonatal sepsis management. In *Jornal de Pediatria* (Vol. 96). Retrieved from DOI: 10.1016/j.jpmed.2019.10.004
- Puopolo KM, Benitz WE, Zaoutis TE. (2018). Management of neonates born at  $\geq 35$  0/7 weeks' gestation with suspected or proven early-onset bacterial sepsis.

*Pediatrics*. 142:e20182894. DOI: 10.1542/peds.2018-2894

- Pusponegoro, T. S. (2016). Sepsis pada Neonatus (Sepsis Neonatal). *Sari Pediatri*, 2(2). Retrieved from <https://doi.org/10.14238/sp2.2.2000.96-102>
- Samsudin, I., & Vasikaran, S. D. (2017). Clinical utility and measurement of procalcitonin. *In Clinical Biochemist Reviews (Vol. 38, Issue 2)*. Retrieved from PMID: 29332972 PMCID: PMC5759088
- Shane AL, Sanchez PJ, Stoll BJ. (2017). Neonatal sepsis. *Lancet*;390:1770–80. DOI: 10.1016/S0140-6736(17)31002-4
- Sharma D, Farahbakhsh N, Shastri S, Sharma P. (2018). Biomarkers for diagnosis of neonatal sepsis: a literature review. *J Matern Fetal Neonatal Med*; 31:1646–59. DOI: 10.1080/14767058.2017.1322060.
- Sproston, N. R., & Ashworth, J. J. (2018). Role of C-reactive protein at sites of inflammation and infection. *In Frontiers in Immunology (Vol. 9, Issue APR)*. DOI: 10.3389/fimmu.2018.00754
- Utomo, M. T., et al. (2021). “Current-proven neonatal sepsis in Indonesian tertiary neonatal intensive care unit: a hematological and microbiological profile.” *Iranian journal of microbiology* vol. 13,; 266-273. doi:10.18502/ijm.v13i3.6386
- Vijayan, A. L., et.al. (2017). Procalcitonin: A promising diagnostic marker for sepsis and antibiotic therapy. *In Journal of Intensive Care (Vol. 5, Issue 1)*. Retrieved from <https://doi.org/10.1186/s40560-017-0246-8>
- Vishnu Tewari, V. (2016). Current Evidence on Prevention and Management of Early Onset Neonatal Sepsis. *Journal of Infectious Diseases & Therapy*, 04(02). doi: 10.4172/2332-0877.1000277
- Wulandari, A., Martuti, S., & Kaswadi, P. (2018). Perkembangan diagnosis sepsis pada anak. *Sari Pediatri*, 19(4). DOI: <http://dx.doi.org/10.14238/sp19.4.2017.237-44>