

DAFTAR PUSTAKA

- American Joint Committee on Cancer. Breast Cancer Staging, ed 7th. Dikutip dari:<https://cancerstaging.org/references/tools/quickreferences/Documents/BreastMedium.pdf>
- Ardina, R., 2018 Respon Inflamasi Pada Perokok Pasif di Kecamatan Pahandut Kota Palangka Raya Ditinjau Dari Jumlah Leukosit dan Jenis Leukosit. *The Journal of Muhammadiyah Medical Laboratory Technology*, pp. Vol. 1 No 2 (31-41). (http://www.researchgate.net/publication/325646468_Respon_Inflamasi_Pada_Perokok_Pasif_Di_Kecamatan_Pahandut_Kota_Palangka_Raya_Ditinjau_Dari_Jumlah_Leukosit_dan_Jenis_Leukosit)
- Argasih, S. 2020. *Hubungan Neutrophil Lymphocyte Ratio (NLR) dan CA 15-3 Dalam Monitoring Prognosis Penderita Kanker Payudara* [online] Diakses dari: <http://repository.unas.ac.id/1715/1/Sartika%20Argasih.pdf> [Pada tanggal 2 November 2022].
- Arulselvan P, Fard MT, Tan WS, Gothai S, Fakurazi S, Norhaizan ME, Kumar SS. Role of Antioxidants and Natural Products in Inflammation. *Oxid Med Cell Longev*. 2016;2016:5276130. doi: 10.1155/2016/5276130. Epub 2016 Oct 10. PMID: 27803762; PMCID: PMC5075620.
- Baratawidjaja, Karnen G. dan Iris Rengganis. 2009. Imunologi dasar. Edisi 8. Jakarta:FK UI
- Bare, Y., Kuki, A.D., Rophi, A.H., Krisnamurti, G.C., Lorenza, M.R.W.G. and Sari, D.R.T. 2019. Prediksi Asam Kuinat Sebagai Anti-Inflamasi Terhadap COX-2 Secara Virtual. *Biota : Jurnal Ilmiah Ilmu-Ilmu Hayati*. 4, 3 (Nov. 2019), 124–129. DOI:<https://doi.org/10.24002/biota.v4i3.2516>.
- Buchari, U. B. 2018. *Perbedaan Kadar CEA Menurut Derajat Histopatologi Adenokarsinoma Rektum di RSUP Haji Adam Malik Medan*. Tesis. Fakultas Kedokteran Universitas Sumatera Utara. Medan
- Candrawati, O., Utomo, B. E. B. H., & Sofii, Imam. (2018). *Correlation of neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio, lymphocyte-to-monocyte ratio and carcinoembryonic antigen level in colorectal cancer*. *JKKI : Jurnal Kedokteran Dan Kesehatan Indonesia*, 9(2), 82–88. <https://doi.org/10.20885/JKKI.Vol9.Iss2.art4>

- Cardoso, F., Kyriakides, S., Ohno, S., Poortmans, P., Rubio, I. T., Zackrisson, S., & Senkus, E. (2019). *Early breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up*. ESMO, 30(8), 1194–1220. <https://doi.org/10.1093/annonc/mdz173>
- Chen W, Liu Q, Tan SY, Jiang YH. Association between carcinoembryonic antigen, carbohydrate antigen 19-9 and body mass index in colorectal cancer patients. *Mol Clin Oncol.* 2013 Sep;1(5):879-886. doi: 10.3892/mco.2013.158. Epub 2013 Jul 23. PMID: 24649265; PMCID: PMC3915429.
- Depkes RI. *Buku Saku Pencegahan Kanker Leher Rahim dan Kanker Payudara*.<https://docs.google.com/file/d/0Bwq8YAw3QBIUkNLbDdSQkxQMWM/edit?pli=1> (situs 25 Juni 2014).
- Duffy MJ. Carcinoembryonic antigen as a marker for colorectal cancer: is it clinically useful? *Clin Chem.* 2001 Apr;47(4):624-30. PMID: 11274010.
- Evan G, Littlewood TA. 1998 matter of life and cell death. (https://www.researchgate.net/publication/13564638_Evan_G_Littlewood_TA_matter_of_life_and_cell_death_Science_281_1317-1322)
- Fallo, I.N., Sidharta, R.A & Gunawan, L.S., 2018. *Korelasi antara Neutrophyl Lympocyte Ratio dengan Stadium Kanker pada Pasien Kanker Payudara. Biomedika*, p Volume. No 02. (<http://ejournal.Setiabudi.ac.id/ojs/index.php/biomedika/article/view/408>)
- Fan S, Xie X, Shen Y, Wang W, Gu X, Yao Z. *The predictive value of preoperative serum neutrophil-to-lymphocyte ratio and tumor markers for early breast cancer patients: A retrospective study*. *Medicine (Baltimore)*. 2022 Aug 12;101(32):e30011. doi: 10.1097/MD.00000000000030011. PMID: 35960055; PMCID: PMC9371529.
- Gago-Dominguez M, Matabuena M, Redondo CM, Patel SP, Carracedo A, Ponte SM, Martínez ME, Castelao JE. *Neutrophil to lymphocyte ratio and breast cancer risk: analysis by subtype and potential interactions*. *Sci Rep.* 2020 Aug 6;10(1):13203. doi: 10.1038/s41598-020-70077-z. Erratum in: *Sci Rep.* 2020 Nov 20;10(1):20641. PMID: 32764699; PMCID: PMC7413522.
- Godinho-mota, J. C. M., Gonçalves, L. V., Mota, J. F., Soares, L. R., Schincaglia, R. M., Martins, K. A., & Freitas-Junior, R. (2019). *Sedentary Behavior and Alcohol Consumption Increase Breast Cancer Risk Regardless of Menopausal Status : A Case-Control Study*. *Nutrients*, 11, 1–9.

- Guthrie GJK, Charles KA, Roxburgh CSD, Horgan PG, McMillan DC, Clarke SJ. *The systemic inflammation-based neutrophil-lymphocyte ratio: experience in patients with cancer.* Crit Rev Oncol Hematol. 2013;88(1):218–30. <http://dx.doi.org/10.1016/j.critrevonc.2013.03.010>.
- Gürağaç A, Demirer Z. The neutrophil-to-lymphocyte ratio in clinical practice. Can Urol Assoc J. 2016 Mar-Apr;10(3-4):141. doi: 10.5489/cuaj.3587. PMID: 27217863; PMCID: PMC4839997.
- Hadjigol S, Shah BA, O'Brien-Simpson NM. The 'Danse Macabre'-Neutrophils the Interactive Partner Affecting Oral Cancer Outcomes. Front Immunol. 2022 Jun 16;13:894021. doi: 10.3389/fimmu.2022.894021. PMID: 35784290; PMCID: PMC9243430.
- Hammarström S. *The carcinoembryonic antigen (CEA) family: structures, suggested functions and expression in normal and malignant tissues.* Semin Cancer Biol. 1999 Apr;9(2):67-81. doi: 10.1006/scbi.1998.0119. PMID: 10202129.
- Handayani, F. W., Muhtadi, A., Farmasi, F., Padjadjaran, U., Dara, T., Manis, K., & Aktif, S. (2013). Aktivitas Anti Kanker Payudara Beberapa Tanaman Herbal. Farmaka, 4, 1–15.
- Han, F., Shang X., Wan F., 2017. *Clinical value of the preoperative neutrophil-to-lymphocyte ratio and red blood cell distribution width in patients with colorectal carcinoma.* Oncology Letters. Edisi 2018: 3339-3342.
- Hermasari, K.B, Ariningrum, D, Subandono J, Mulyani S, Hastuti H.,2019. *Buku Pedoman Keterampilan Klinis. Teknik Injeksi dan Pungsi.* Laboratorium Keterampilan Klinis. Surakarta: Fakultas Kedokteran Sebelas Maret Surakarta (<https://skillslab.fk.uns.ac.id/wp-content/uploads/2019/02/smt-4-KETERAMPILAN-INJEKSI-DAN-PUNGSI-2019.pdf>)
- Huang ZB, Zhou X, Xu J, Du YP, Zhu W, Wang J, Shu YQ, Liu P. Prognostic value of preoperative serum tumor markers in gastric cancer. World J Clin Oncol. 2014 May 10;5(2):170-6. doi: 10.5306/wjco.v5.i2.170. PMID: 24829865; PMCID: PMC4014790.
- Imamura M, Morimoto T, Egawa C, Fukui R, Bun A, Ozawa H, Miyagawa Y, Fujimoto Y, Higuchi T, Miyoshi Y. Significance of baseline neutrophil-to-lymphocyte ratio for progression-free survival of patients with HER2-positive breast cancer treated with trastuzumab emtansine. Sci Rep. 2019 Feb 12;9(1):1811. doi: 10.1038/s41598-018-37633-0. PMID: 30755651; PMCID: PMC6372578.

- Iqmy, L. O., Setiawati, & Yanti, D. E. (2021). *FAKTOR RISIKO YANG BERHUBUNGAN DENGAN KANKER PAYUDARA*. Jurnal Kebidanan, 7(1), 32–36.
- Kalli, S., Semine, A., Cohen, S., Naber, S. P., Makim, S. S., & Bahl, M. (2018). *American Joint Committee on Cancer's Staging System for Breast Cancer*. RadioGraphics, 38(7), 1921–1933.
- Kemenkes RI. (2011). Pedoman Interpretasi Data Klinik.
- Kemenkes RI, 2018 Pedoman Nasional Pelayanan Kedokteran Tata Laksana Kanker Payudara .
https://yankes.kemkes.go.id/unduhan/fileunduhan_1610414392_632733.pdf
- KEMENPPPA, 2022. *KEMENPPPA DORONG MASYARAKAT LAKUKAN DETEKSI DINI KANKER PAYUDARA DENGAN SADARI DAN SADANIS*. www.kemenpppa.go.id
- Khasanah, 2013. Karsinoma Mammea Stadium IV Dengan Tanda-Tanda Dyspnoe Dan Paraplegi Ekstremitas Inferior. Medula, Volume 1, Nomor 2, Oktober 2013. Lampung
- Kumar, V. Abbas, A.K, Aster, J.C. (2018). Robins *Basic Pathology Edisi ke-10*. Philadelphia: Elsevier
- Lee EY, Muller WJ. Oncogenes and tumor suppressor genes. *Cold Spring Harb Perspect Biol*. 2010 Oct;2(10):a003236. doi: 10.1101/csdperspect.a003236. Epub 2010 Aug 18. PMID: 20719876; PMCID: PMC2944361.
- Li Y, Cao H, Jiao Z, Pakala SB, Sirigiri DN, Li W, Kumar R, Mishra L. Carcinoembryonic antigen interacts with TGF-{beta} receptor and inhibits TGF-{beta} signaling in colorectal cancers. *Cancer Res*. 2010 Oct 15;70(20):8159-68. doi: 10.1158/0008-5472.CAN-10-1073. Epub 2010 Oct 1. PMID: 20889724; PMCID: PMC3001246.
- Locker GY, Hamilton S, Harris J, Jessup JM, Kemeny N, Macdonald JS, Somerfield MR, Hayes DF, Bast RC Jr; ASCO. *ASCO 2006 update of recommendations for the use of tumor markers in gastrointestinal cancer*. *J Clin Oncol*. 2006 Nov;24(33):5313-27. doi: 10.1200/JCO.2006.08.2644. Epub 2006 Oct 23. PMID: 17060676.
- Made. 2014. *Kanker dan Antikanker*. Bahan Ajar. Universitas Udayana

Markiewski MM, Lambris JD. The role of complement in inflammatory diseases from behind the scenes into the spotlight. *Am J Pathol.* 2007 Sep;171(3):715-27. doi: 10.2353/ajpath.2007.070166. Epub 2007 Jul 19. PMID: 17640961; PMCID: PMC1959484.

Master degree; Shen, Yong PhD; Wan, Wenjun (2022). *The Predictive value of preoperative serum neutrophil-to-lymphocyte ratio and tumor markers for early breast cancer patients: A retrospective study.* Medicine 101(32):p e300311

Mindray, 2019. BC-6800 Operator's Manual V10.0 en. (<https://www.scribd.com/document/419832333/BC-6800-Operator-s-Manual-V10-0-En>)

Momenimovahed, Z., & Salehiniya, H. (2019). *Epidemiological characteristics of and risk factors for breast cancer in the world.* Dovepress, 11, 151–164.

Moo TA, Sanford R, Dang C, Morrow M. Overview of Breast Cancer Therapy. *PET Clin.* 2018 Jul;13(3):339-354. doi: 10.1016/j.cpet.2018.02.006. PMID: 30100074; PMCID: PMC6092031.

Nagpal M, Singh S, Singh P, Chauhan P, Zaidi MA. *Tumor markers: A diagnostic tool.* *Natl J Maxillofac Surg.* 2016 Jan-Jun;7(1):17-20. doi: 10.4103/0975-5950.196135. PMID: 28163473; PMCID: PMC5242068.

Nicholson BD, Shinkins B, Pathiraja I, Roberts NW, James TJ, Mallett S, Perera R, Primrose JN, Mant D. Blood CEA levels for detecting recurrent colorectal cancer. *Cochrane Database Syst Rev.* 2015 Dec 10;2015(12):CD011134. doi: 10.1002/14651858.CD011134.pub2. PMID: 26661580; PMCID: PMC7092609.

Noh H, Eomm M, Han A. *Usefulness of pretreatment neutrophil to lymphocyte ratio in predicting disease-specific survival in breast cancer patients.* *J Breast Cancer.* 2013;16(1):55–9. <http://dx.doi.org/10.4048/jbc.2013.16.1.55>.

Nurrohmah, A., Aprianti, A., & Hartutik, S. (2022). *Risk Factors of Breast Cancer.* *GASTER JOURNAL OF HEALTH SCIENCE*, 20(1), 1–10.

O. Orakpoghenor, D. O. Avazi, T. P. Markus, and O. S. Olaolu, “*Lymphocytes : A Brief Review -*,” vol. 3, no. June, pp. 4–8, 2019.

Pangribowo, S. (2019). *Beban Kanker di Indonesia. Pusat Data Dan Informasi Kementerian Kesehatan RI*, 1–16.

Panigoro, S., Hernowo, BS., Purwanto, H ., 2015. *Panduan Penatalaksanaan Kanker Payudara* . Jakarta Komite Penanggulangan Kanker Nasional, Kemenkes R1. (<http://kanker.kemkes.go.id/guidelines/PPKPayudara.pdf>)

Pedrazzani C, Mantovani G, Fernandes E, Bagante F, Luca Salvagno G, Surci N, Campagnaro T, Ruzzenente A, Danese E, Lippi G, Guglielmi A. *Assessment of neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio and platelet count as predictors of long-term outcome after R0 resection for colorectal cancer.* Sci Rep. 2017 May 4;7(1):1494. doi: 10.1038/s41598-017-01652-0. PMID: 28473700; PMCID: PMC5431463.

Prasetyo, Y.E., Bahrun, U & Pakasi, R.D., 2015. *Angka Banding Neutrofil/Limfosit di Karsinoma Payudara. Indonesia Journal of Clinical Pathology and Medical Laboratory*, pp.125-129. (<http://indonesianjounalofclinicalpathhology.org/article/view/1090/0>)

Pulungan, R.M., 2010. Karakteristik Penderita Kanker Payudara Rawat Inap di RS Haji Medan Tahun 2005-2009. Skripsi. FKM USU Medan.

Puspitawati, D. A. (2018). *SISTEM PAKAR DIAGNOSIS PENYAKIT KANKER PAYUDARA DAN CARA PENANGANANNYA*. Jurnal TECHNO Nusa Mandiri, 15(2), 129–136.

Raehaan, N.R., Nurulita, A. and Arif, M. 2016. *CARCInoeMBRYonIC AnTIGen (CEA) DI KANKER KOLOREKTAL. INDONESIAN JOURNAL OF CLINICAL PATHOLOGY AND MEDICAL LABORATORY*. 20, 3 (Oct. 2016), 192–196. DOI:<https://doi.org/10.24293/ijcpml.v20i3.465>.

Rahfiludin, M. Z., Dina, R.A., 2016. *Dasar imunologi Bagi Kesehatan Masyarakat*. Yogyakarta: Trans Medika.

Rahma Isti Kusyadi, Florentina Sustini, Eddy Herman Tanggo. 2014. *Profil Cea Dan Ca 15-3 Pasien Kanker Payudara Di Poli Bedah Onkologi Rsud Dr.Soetomo Surabaya*. Fakultas Kedokteran, Universitas Airlangga, Surabaya, Jawa Timur

Reddy JP, Hernandez M, Gunther JR, Dabaja BS, Martin GV, Jiang W, Akhtari M, Allen PK, Atkinson BJ, Smith GL, Pinnix CC, Milgrom SA, Abou Yehia Z, Osborne EM, Oki Y, Lee H, Hagemeister F, Fanale MA. Pre-treatment neutrophil/lymphocyte ratio and platelet/lymphocyte ratio are prognostic of progression in early stage classical Hodgkin lymphoma. Br J Haematol. 2018 Feb;180(4):545-549. doi: 10.1111/bjh.15054. Epub 2017 Dec 21. PMID: 29271057.

- Retno MW, Suci DR, Rizana F, 2019. *Carcinoembryonic Antigen (CEA) dan Neutrofilo-Limfosit Ratio (NLR) sebagai Faktor Prediktif Kanker Kolorektal*. Ilmu Teknologi Kesehatan , Vol. 7, Nomor 1, September 2019, hlm:68-76
- Roche Diagnostics, 2022. Cobas e411 analyzer. (<https://diagnostics.roche.com/global/en/products/instruments/cobas-e-411-ins-502.html>)
- Romadhon, Y. A. (2013). Gangguan Siklus Sel dan Mutasi Gen pada Kanker Payudara. 40(10), 786–789. Retrieved from http://www.kalbemed.com/Portals/6/1_25_209Opini-Gangguan Siklus Sel dan Mutasi Gen pada Kanker Payudara.pdf
- Sacher, A. R., 2004. *Tinjauan Klinis Hasil Pemeriksaan Laboratorium*. Jakarta: Gramedia Pustaka Umum.
- Savitri, Astrid. 2015. *Kupas Tuntas Kanker Payudara, Leher Rahim, dan Rahim*. Yogyakarta, Pustaka Baru Press.
- Setiawan, S. D., 2015. The Effect of Chemotherapy in Cancer Patient to Anxiety. J MAJORITY, p. Volume 4 Nomor 4. (<http://juke.kedokteran.unila.ac.id/index.php/majority/article/view/587>)
- Sukowati, E.G. 2011. *Pengaruh Pemberian Eicosapentaeonic Acid (EPA) terhadap Jumlah Sel T CD4 pada Pasien Karsinoma Mamae Stadium III yang Mendapatkan Kemoterapi* [online] Diakses dari:<http://eprints.undip.ac.id/33602/>[Pada tanggal 10 Mei 2013].
- Templeton AJ, McNamara MG, Seruga B, Vera-Badillo FE, Aneja P, Ocana A, Leibowitz-Amit R, Sonpavde G, Knox JJ, Tran B, Tannock IF, Amir E. *Prognostic role of neutrophil-to-lymphocyte ratio in solid tumors: a systematic review and meta-analysis*. J Natl Cancer Inst. 2014;106(6):dju124. doi:10.1093/jnci/dju124.
- Treffers LW, Hiemstra IH, Kuijpers TW, van den Berg TK, Matlung HL. *Neutrophils in cancer*. Immunol Rev. 2016 Sep;273(1):312-28. doi: 10.1111/imr.12444. PMID: 27558343.
- Tsai, P., Su, W., Leung, W., 2016. *Neutrophil–lymphocyte ratio and CEA level as prognostic and predictive factors in colorectal cancer: A systematic review and meta-analysis*. Journal of Cancer Research and Therapeutics. Edisi April - Juni 2016. Vol 12 No 2: 582-587.

- Ulas A, Avci N, Kos T, Cubukcu E, Olmez OF, Bulut N, Degirmenci M. Are neutrophil/lymphocyte ratio and platelet/lymphocyte ratio associated with prognosis in patients with HER2-positive early breast cancer receiving adjuvant trastuzumab? *J BUON.* 2015 May-Jun;20(3):714-22. PMID: 26214622.
- Wang JY, Lu CY, Chu KS, Ma CJ, Wu DC, Tsai HL, Yu FJ, Hsieh JS. *Prognostic significance of pre- and postoperative serum carcinoembryonic antigen levels in patients with colorectal cancer.* Eur Surg Res. 2007;39(4):245-50. doi: 10.1159/000101952. Epub 2007 Apr 23. PMID: 17457032.
- Xiao J, Ye ZS, Wei SH, Zeng Y, Lin ZM, Wang Y, Teng WH, Chen LC. Prognostic significance of pretreatment serum carcinoembryonic antigen levels in gastric cancer with pathological lymph node-negative: A large sample single-center retrospective study. *World J Gastroenterol.* 2017 Dec 28;23(48):8562-8569. doi: 10.3748/wjg.v23.i48.8562. PMID: 29358864; PMCID: PMC5752716.
- Yin X, Xiao Y, Li F, Qi S, Yin Z, Gao J. Prognostic Role of Neutrophil-to-Lymphocyte Ratio in Prostate Cancer: A Systematic Review and Meta-analysis. *Medicine (Baltimore).* 2016 Jan;95(3):e2544. doi: 10.1097/MD.0000000000002544. PMID: 26817900; PMCID: PMC4998274.
- Zhao W, Li X, Wang W, Chen B, Wang L, Zhang N, Wang Z, Yang Q. Association of Preoperative Serum Levels of CEA and CA15-3 with Molecular Subtypes of Breast Cancer. *Dis Markers.* 2021 Sep 27;2021:5529106. doi: 10.1155/2021/5529106. PMID: 34621407; PMCID: PMC8492280.